

**Black-capped Chickadee**



**Boreal Chickadee**



*Chickadees are common, spunky feeder visitors. ID using black vs. brown cap.*



Observer: \_\_\_\_\_

Mailing Address: \_\_\_\_\_

Email: \_\_\_\_\_ Add to ASI email list? \_\_\_\_\_

Section of Borough: \_\_\_\_\_

Enclosed is my gift of \$ \_\_\_\_\_ to support Fairbanks FeederCount

**Number of Birds at your Feeders (see instructions)**

Winter 2025-26	Nov 8	Dec 6	Mar 7
Ruffed Grouse			
Spruce Grouse			
Rock Dove (Pigeon)			
Downy Woodpecker			
Hairy Woodpecker			
Canada Jay			
Black-capped Chickadee			
Boreal Chickadee			
Red-breasted Nuthatch			
Pine Grosbeak			
Redpolls			
Dark-eyed Junco			
Other:			
Watched but none seen			
Temperature/Weather			

**Redpoll**




*Grosbeaks & redpolls are finches. They often visit feeders in groups. Male redpolls' chests turn reddish in spring. Note: The American Ornithological Society now considers Common and Hoary Repolls to be a single species, known simply as Repoll.*



**Pine Grosbeak**

**Downy Woodpecker**



**Canada Jay**

*Downy Woodpeckers are smaller than Hairy Woodpeckers. Also note the smaller bill (relative to head size).*



**Red-breasted Nuthatch**



**Hairy Woodpecker**



**Dark-eyed Junco**

*Watch for late migratory species like juncos & robins.*

Illustrations generously provided by George West [www.birchsidestudios.com](http://www.birchsidestudios.com)

**Types of Food (total quantity)**

Black oil sunflower seeds
Sunflower chips
Mixed seed
Thistle (niger)
Suet (fat)
Peanut butter
Grit

**Types of Feeders (# of each)**

Tray or Shelf
Hopper/enclosed
Tube
Clinger
Hanging Suet
Suet on tree

**Outdoor Pets (# of each)**

Cats
Dogs

**Housing Density (check one)**

High (≤ 0.5 acre lots)
Moderate (0.6-2 acres)
Low (2-5 acre lots)
Rural (< 1 house/5 acres)

**Habitat (approximate % of each)**

River/Stream
Wetland
Shrub* (≤ 2.5 m)
Mixed Forest*
Deciduous Forest*
Coniferous Forest*

\*Please list dominant species if possible.

